arXiv:hep-ph/0111281v1 22 Nov 2001

On the Mixed Propagator Approach to $\rho-\omega$ Mixing *

Mu-Lin Yan†, Ji-Hao Jiang
CCST(World Lab), P.O.Box 8730, Beijing,100080, P.R.China
and
Center for Fundamental Physics,University of Science and Technology of China‡
Hefei, Anhui 230026, China

Xiao-Jun Wang
Institution of Theoretical Physics, Beijing, 100080, P.R.China

Mixed Propagator (MP) approach to $\rho-\omega$ mixing is discussed. It is found that under the pole-approximation assumption the results of MP approach is not compatible both with the effective Lagrangian theory and with the experiment measurement criterion. To overcome these inconsistent, we propose a new MP approach in which the physical states of ρ and ω are determined by the requirement of experimental measurement to meson resonance. In terms of this new MP approach, the EM pion form factor F_π and form factors of $\rho^0\to\pi^0\gamma$ and of $\omega\to\pi^0\gamma$ are derived. The results of F_π are in good agreement with data. The form factor of $\rho^0\to\pi^0\gamma$ exhibits a hidden charge-asymmetry enhancement effect which agree with the prediction of the effective Lagrangian theory.

14.40.Cs,13.25.Jx,12.40.Vv,13.40.Hq

$\rho^0-\omega$ mixing (or $\rho^0-\omega$ interference, or interference between overlapping resonances, or charge asymmetry caused by $\rho-\omega$ mixing and so on) has attracted much interest during past four decades. Since 1961 [1], there is a great deal of works on this subject in the literature (see, for example, review articles [2,3] and the references within). Recently, we used model-independent effective Lagrangian (eff-L) approach to discuss $\rho-$ and $\omega-$ anomaly-like processes of $\rho^0\to\pi^0\gamma$, and of $\omega\to\pi^0\gamma$ [4]. It has been found that there exist a hidden charge-asymmetry enhancement effect to $\rho^0\to\pi^0\gamma$ due to $\rho^0-\omega$ mixing. Namely, we have (by using the notations in ref [4])

$$\begin{aligned} f_{\rho^0\pi^0\gamma}|_{\rm eff-L} &= f^{(0)}_{\rho^0\pi^0\gamma}\left(1+\frac{3\Pi_{\rho\omega}(m_\rho^2)}{m_\rho^2-m_\omega^2+im_\omega\Gamma_\omega}\right), \\ f_{\omega\pi^0\gamma}|_{\rm eff-L} &= f^{(0)}_{\omega\pi^0\gamma}\left(1+\frac{1}{3}\frac{\Pi_{\rho\omega}(m_\omega^2)}{m_\omega^2-m_\rho^2+im_\rho\Gamma_\rho}\right), \end{aligned} \tag{1}$$

where the subscript 'eff-L' means the results obtained by in the effective Lagrangian approach. Due to $\Gamma_\omega << \Gamma_\rho$, we have

$$|f_{\rho^0\pi^0\gamma}|_{\rm eff-L} >> f^{(0)}_{\rho^0\pi^0\gamma} = f_{\rho^\pm\pi^\pm\gamma} \tag{2}$$

$$|f_{\omega\pi^0\gamma}|_{\rm eff-L} \sim f^{(0)}_{\omega\pi^0\gamma}. \tag{3}$$

This reflects an unusual charge asymmetry enhancement effect for $\rho\to\pi\gamma$. As addressed in [4], this hidden effect has already been implied by experiment [5].

On the other hand, There is a well-known quantum mechanics method to deal with the $\rho-\omega$ mixing problem: the approach of mixed propagator with pole approximation (shortly, we will call it as pole-Mixed-Propagator approach, or pole-MP approach) [2,3]. This approach was developed even before discovery of QCD. A question arisen here what are results for $\rho^0\to\pi^0\gamma$ and $\omega\to\pi^0\gamma$ in pole-MP approach? The vector meson propagator is given by (Renard representation)

---

† mlyan@ustc.edu.cn

* This work is partly supported by the grant No.90103002 of National Natural Science Foundation of China.

‡ Mailing address

$$
\begin{aligned}
D_{\mu\nu}(q^2) &= \int d^4x e^{-iq.x}\langle 0|T\{V_\mu(x)V_\nu(0)\}|0\rangle \\
&= D(q^2)g_{\mu\nu} + \frac{1}{q^2}(D(0)-D(s))q_\mu q_\nu
\end{aligned}
\tag{4}
$$

where $s \equiv q^2$, and the propagator function $D(s)$ is written in the following way

$$
D(s) = \frac{1}{s - W(s)}.
\tag{5}
$$

For multi-vector-meson channels, $W(s)$ is complex mass-square matrix with non-zero off-diagonal elements in general. The basic assumption for pole-MP approach is that the physical states are eigenvectors of $W(s)$, or the complex mass-square of physical state is determined by the following equation

$$
det[s - W(s)] = 0.
\tag{6}
$$

For $\rho - \omega$ case, the solutions of this physical condition equation are follows [3]

$$
\begin{aligned}
|\rho^0_p\rangle &= |\rho^0_I\rangle + \epsilon|\omega_I\rangle, \quad \langle\tilde{\rho}^0_p| = \langle\rho^0_I| - \epsilon\langle\omega_I| \\
|\omega_p\rangle &= |\omega_I\rangle - \epsilon|\rho^0_I\rangle, \quad \langle\tilde{\omega}_p| = \langle\omega^0_I| + \epsilon\langle\rho^0_I|
\end{aligned}
\tag{7}
$$

with

$$
\epsilon = \frac{\Pi_{\rho\omega}}{m^2_\omega - m^2_\rho + i(m_\omega\Gamma_\omega - m_\rho\Gamma_\rho)},
\tag{8}
$$

where $|\rho^0_p\rangle$ and $|\omega_p\rangle$ denote physical ρ^0- and $\omega-$ states under the pole approximation assumption respectively, $|\rho^0_I\rangle \equiv |1,0\rangle$ and $|\omega_I\rangle \equiv |0,0\rangle$ are in the isospin basis, and $\Pi_{\rho\omega} \equiv \langle\rho^0_I|W|\omega_I\rangle$. Note that ϵ is not a real number, and hence the transfer matrix from isospin basis to physical basis is not unitary. The physical states are normalized in following way

$$
\langle\tilde{a}|b\rangle = \delta_{ab} \quad \text{and} \quad \mathbf{I} = \sum_a |\tilde{a}\rangle\langle a| = \sum_a |a\rangle\langle\tilde{a}|
\tag{9}
$$

where $a,\ b = \rho^0_p$ or ω_p. Because the transfer matrix between the isospin bases and the physical one is not unitary in the pole-MP, the relations of eq. (7) can not be realized in the formalism of unitary Lagrangian field theory. Since it has been widely accepted that the effective Lagrangian field theories inspired by QCD conceptions (e.g., VMD, large N_c expansion, non-linear realization of chiral symmetry, WCCWZ realization, WZ-anomaly, hidden local symmetry and so on) is legitimate and useful to study $\rho-$ and $\omega-$ physics [6], that the pole-MP approach contradicts to effective Lagrangian field theory is a difficulty for pole-MP.

To show the differences between two approach further, let us consider $\rho^0, \omega \to \pi^0\gamma$ decays in the pole-MP. Again, using $f_{\rho^0\pi^0\gamma}$, $f_{\omega\pi^0\gamma}$ and $f^{(0)}_{\rho^0\pi^0\gamma}$, $f^{(0)}_{\omega\pi^0\gamma}$ to denote the physical coupling constants and the isospin-basis constants respectively, we have

$$
\begin{aligned}
f_{\rho^0\pi^0\gamma}|_{\text{pole}-\text{MP}} &= f^{(0)}_{\rho^0\pi^0\gamma}\left(1 + \frac{3\Pi_{\rho\omega}(m^2_\rho)}{m^2_\rho - m^2_\omega + i(m_\omega\Gamma_\omega - m_\rho\Gamma_\rho)}\right), \\
f_{\omega\pi^0\gamma}|_{\text{pole}-\text{MP}} &= f^{(0)}_{\omega\pi^0\gamma}\left(1 + \frac{1}{3}\frac{\Pi_{\rho\omega}(m^2_\omega)}{m^2_\omega - m^2_\rho - i(m_\omega\Gamma_\omega - m_\rho\Gamma_\rho)}\right).
\end{aligned}
\tag{10}
$$

where $f^{(0)}_{\omega\pi^0\gamma} = 3f^{(0)}_{\rho^0\pi^0\gamma}$ has been used. Comparing eq. (1) with eq. (10) and noting $\Gamma_\omega << \Gamma_\rho$, we can see

$$
\begin{aligned}
f_{\rho^0\pi^0\gamma}|_{\text{pole}-\text{MP}} &\sim f^{(0)}_{\rho^0\pi^0\gamma} << f_{\rho^0\pi^0\gamma}|_{\text{eff}-\text{L}} \\
f_{\omega\pi^0\gamma}|_{\text{pole}-\text{MP}} &\sim f^{(0)}_{\omega\pi^0\gamma} \sim f_{\omega\pi^0\gamma}|_{\text{eff}-\text{L}}
\end{aligned}
\tag{11}
$$

Eq. (11) indicates there is no the hidden charge-asymmetry enhancement effect for $\rho^0 \to \pi^0\gamma$ mentioned above in the pole-MP formalism. The result of pole-MP approach to $\rho^0 \to \pi^0\gamma$ is significantly different from one of eff-L. Thus we learned that there may be something wrong. The purpose of this letter is to re-examine the pole-Mixed Propagator approach, and propose a new mixed propagator approach based on the criterion of experimental measurement to meson resonances.

Let's consider whether the pole-MP assumption on physical states eq. (6) is consistent with the experimental criterion to measure the physical meson resonance or not. We first consider the single vector meson resonance channel case. The $D_{\mu\nu}$ of eq. (4) is now the ordinary propagator of a vector meson. The reaction amplitude for a process by the medium of this vector meson resonance reads

$$\mathcal{M} \sim J_1^{\mu} D_{\mu\nu} J_2^{\nu} = (J_1 \cdot J_2) \frac{1}{s - W(s)} \tag{12}$$

where J_1^{μ} and J_2^{ν} represent some currents and $q_\mu J^{\mu}_{1,\,2} = 0$. The reaction probability is [8]

$$\sigma(s) \sim |\mathcal{M}|^2 \propto |\frac{1}{s - W(s)}|^2 \tag{13}$$

The meson resonance mass measured in the experiment is real and is determined by the location of the maximum of $\sigma(s)-$peak in real $s-$axis. Then the resonance mass is determined by the following equation

$$\frac{\partial}{\partial s} |s - W(s)|^2 = 0 \tag{14}$$

Consequently, the mass-square of the resonance, M^2, is determined by the solution of above equation, i.e.,

$$s = [ReW - \frac{1}{4}\frac{\partial}{\partial s}(W^* W + W W^*)](1 - \frac{\partial}{\partial s} ReW)^{-1} \equiv M^2(s). \tag{15}$$

Now return to $\rho^0 - \omega$ two channel case, $W(s)$ and hence $M^2(s)$ are 2×2 matrices. The mass determination equation for physical resonance states should read

$$det[s - M^2(s)] = 0. \tag{16}$$

The physical states then are the eigenvectors of the real mass-matrix M^2. Obviously, the corresponding pole-MP equation (6) is definitely different from above mass determination equation derived from the experiment measurement principle to physical resonance states. It indicates that the the pole-MP is not consistent with the criterion of experimental measurement to the physical resonances. Consequently, we should use eq. (16) to define true physical $\rho-$ and $\omega-$fields, and rule out the condition of eq. (6). For convenience, we will call this method hereafter as Experiment-criterion Mixed Propagator approach, or expt-MP approach.

Now let us use expt-MP to fit experiments. Following refs [2,3] and using Breit-Wigner approximation, we have

$$W = \begin{pmatrix} m^2_{\rho_I} - i\sqrt{s}\Gamma_{\rho_I} & \Pi_{\rho\omega}(s) \\ \Pi_{\rho\omega}(s) & m^2_{\omega_I} - i\sqrt{s}\Gamma_{\omega_I} \end{pmatrix}$$

Considering Im$\Pi_{\rho\omega}$ is small and hence ignorable [9], we get

$$ReW = \begin{pmatrix} m^2_{\rho_I} & \Pi_{\rho\omega}(s) \\ \Pi_{\rho\omega}(s) & m^2_{\omega_I} \end{pmatrix}$$

Then, in $\partial ReW/\partial s$, only off-diagonal elements of $\partial \Pi_{\rho\omega}/\partial s$ is left. Generally, for a broad class of models $\Pi_{\rho\omega}(s)$ at $(\rho-, \omega-)$resonance energy region can be determined by taking VMD-type $\rho - \omega$ mixing Lagrangian $\mathcal{L}_{\rho\omega} = f_{\rho\omega} \rho^{\mu\nu} \omega_{\mu\nu}$ ($V^{\mu\nu} = \partial^\mu V^\nu - \partial^\nu V^\mu,\ V = \rho, \omega$). It leads to $\Pi_{\rho\omega}(s) = f_{\rho\omega} s$ which satisfies $\Pi(s = 0)_{\rho\omega} = 0$ required by generic consideration in ref [10]. Thus, we have

$$\frac{\partial}{\partial s}\Pi_{\rho\omega} = f_{\rho\omega} = \frac{\Pi_{\rho\omega}}{s}|_{s \sim m^2_\rho} \simeq 6.7 \times 10^{-3} \ll 1 \implies 1 - (\partial ReW/\partial s) \simeq 1. \tag{17}$$

where $|\Pi_{\rho\omega}| \simeq 4000 MeV^2$ [10] has been used in the estimation. Furthermore, noting $\Gamma_\omega / m_\omega \ll 1$, we have

$$M^2 = \begin{pmatrix} m^2_{\rho_I} - \frac{\partial(s\Gamma^2_\rho)}{\partial s} & \Pi_{\rho\omega} - \frac{1}{2}(m^2_{\rho_I} + m^2_{\omega_I})\frac{\partial}{\partial s}\Pi_{\rho\omega} \\ \Pi_{\rho\omega} - \frac{1}{2}(m^2_{\rho_I} + m^2_{\omega_I})\frac{\partial}{\partial s}\Pi_{\rho\omega} & m^2_{\omega_I} \end{pmatrix} \tag{18}$$

where the off-diagonal elements of M^2 matrix represent the $\rho-\omega$ mixing in the isospin basis. The physical ρ and ω are eigenstates of M^2. Because M^2 matrix is real and symmetric, distinguishing from the transformation from isospin basis to physical basis in the pole-MP, the corresponding transfer-matrix of expt-MP is unitary. M^2 matrix is diagonalized by the unitary 2×2 matrix C:

$$CM^2C^\dagger = \begin{pmatrix} m_\rho^2 & 0 \\ 0 & m_\omega^2 \end{pmatrix}$$

where

$$C = \begin{pmatrix} 1 & -\eta \\ \eta & 1 \end{pmatrix}, \qquad \eta = -\frac{\Pi_{\rho\omega}(1-\frac{1}{2s}(m_\rho^2+m_\omega^2))}{(m_\omega^2-m_\rho^2)}. \tag{19}$$

Consequently, the solutions of expt-MP's physical state condition of eq. (16) are follows

$$\begin{aligned} |\rho_p^0\rangle &= |\rho_I^0\rangle - \eta|\omega_I\rangle, \;\; \langle\rho_p^0| = |\rho_p^0\rangle^\dagger \\ |\omega_p\rangle &= |\omega_I\rangle + \eta|\rho_I^0\rangle, \;\; \langle\omega_p| = |\omega_p\rangle^\dagger. \end{aligned} \tag{20}$$

Under this transformation, we have

$$CWC^\dagger = \begin{pmatrix} z_\rho & T \\ T & z_\omega \end{pmatrix} \tag{21}$$

where $z_\rho = m_\rho^2 - im_\rho\Gamma_\rho, z_\omega = m_\omega^2 - im_\omega\Gamma_\omega$, and $T = \Pi_{\rho\omega} - \eta(z_\omega - z_\rho)$ are all defined in the physical state basis of expt-MP. The propagator function in the physical basis D^P reads

$$\begin{aligned} D^P(s) &= C(s-W)^{-1}C^\dagger = (s-CWC^\dagger)^{-1} \\ &= \begin{pmatrix} (s-z_\rho)^{-1} & (s-z_\rho)^{-1}T(s-z_\omega)^{-1} \\ (s-z_\omega)^{-1}T(s-z_\rho)^{-1} & (s-z_\omega)^{-1} \end{pmatrix} \\ &\equiv \begin{pmatrix} D^P_{\rho\rho} & D^P_{\rho\rho}TD^P_{\omega\omega} \\ D^P_{\omega\omega}TD^P_{\rho\rho} & D^P_{\omega\omega} \end{pmatrix} \equiv \begin{pmatrix} D^P_{\rho\rho} & D^P_{\rho\omega} \\ D^P_{\omega\rho} & D^P_{\omega\omega} \end{pmatrix}. \end{aligned} \tag{22}$$

For $V-X$-vertex ($V=\rho,\ \omega$ and X represents other particles), f^F_{VX} denotes the corresponding form-factor, f^P_{VX} and f^0_{VX} denote the coupling constants in the physical basis and in the isospin basis respectively.



FIG. 1. The relation between the form factor of $V-X$ vertex and the corresponding coupling constants for the vertices, where $V=\rho$ in Fig.(a) and $V=\omega$ in Fig.(b). In the left-hand side, the black dots in the vertex represent the form factor. In the right-hand side, the coupling constants are at the vertices. The single thick (dash) lines denote for $\rho-$propagator $D^P_{\rho\rho}$ ($\omega-$propagator $D^P_{\omega\omega}$), and the thick-cross-dash lines (or dash-cross-thick) line denote the mixed propagator $D^P_{\rho\omega}$ (or $D^P_{\omega\rho}$). The thin lines are external lines of X-particles.

Since $D^P_{\rho\omega} = D^P_{\omega\rho} \neq 0$ in the expt-MP, generally, the form factor is different from the corresponding coupling constant, i.e., $f^F_{VX} \neq f^P_{VX}$ (by contrast, $f^F_{VX} = f^P_{VX}$ in pole-MP approach due to $D^P_{\rho\omega} = D^P_{\omega\rho} = 0$). From Fig.1, we have

$$D^P_{VV}f^F_{VX} = D^P_{V\rho}f^P_{\rho X} + D^P_{V\omega}f^P_{\omega X}, \tag{23}$$

with

$$f^P_{\rho X} = f^{(0)}_{\rho X} - \eta f^{(0)}_{\omega X} \qquad\qquad f^P_{\omega X} = f^{(0)}_{\omega X} + \eta f^{(0)}_{\rho\gamma}. \tag{24}$$

In terms of eqs. (22) and (24) the time-like EM pion form-factor is given, in the $\rho - \omega$ interference region, by

$$\begin{aligned} F_\pi(s) &= 1 + [f^P_{\rho\gamma} D^P_{\rho\rho} f^F_{\rho\pi\pi} + f^P_{\omega\gamma} D^P_{\omega\omega} f^F_{\omega\pi\pi}] \\ &= 1 + \frac{f^P_{\rho\gamma} f^P_{\rho\pi\pi}}{s - z_\rho} + \frac{f^P_{\omega\gamma} f^P_{\omega\pi\pi}}{s - z_\omega} + \frac{(f^P_{\rho\gamma} f^P_{\omega\pi\pi} + f^P_{\omega\gamma} f^P_{\rho\pi\pi})T}{(s - z_\rho)(s - z_\omega)} \\ &= 1 + (f^P_{\rho\gamma} f^P_{\rho\pi\pi} + \frac{(f^P_{\rho\gamma} f^P_{\omega\pi\pi} + f^P_{\omega\gamma} f^P_{\rho\pi\pi})T}{z_\rho - z_\omega})(\frac{1}{s - z_\rho} + \xi e^{i\phi} \frac{1}{s - z_\omega}), \end{aligned} \tag{25}$$

with

$$\xi e^{i\phi} = [\frac{1}{3}\eta - \frac{(\eta + \frac{1}{3})T}{z_\rho - z_\omega}][1 + \frac{(\eta + \frac{1}{3})T}{z_\rho - z_\omega}]^{-1}, \tag{26}$$

where $f^{(0)}_{\rho\gamma} = 3 f^{(0)}_{\omega\gamma}$ and $f^{(0)}_{\omega\pi\pi} = 0$ have been used, and ϕ is Orsay phase. Using $\Pi_{\rho\omega} \simeq -4000 MeV^2$ [10] in eq. (26), we obtain that $\xi \simeq 0.012$ and ϕ is equal to about $100^o \to 101^o$ as s varies from m^2_ρ to m^2_ω. These predictions are in good agreement with experimental data [11], and hence the expt-MP approach is legitimate to describe the $\rho - \omega$ mixing effects in the pion EM-form factor.



FIG. 2. The Electromagnetic(EM) pion form factor. Indications of the lines for the propagators are same as Fig.1. The black dots are form factors.

Now, we study the anomalous-like $\rho^0 \to \pi^0\gamma$ and $\omega \to \pi^0\gamma$ decays in terms of the expt-MP approach. Namely, taking $X = \pi^0\gamma$ in eq. (23), we have

$$\begin{aligned} D^P_{\rho\rho} f^F_{\rho^0\pi^0\gamma} &= D^P_{\rho\rho} f^P_{\rho^0\pi^0\gamma} + D^P_{\rho\omega} f^P_{\omega\pi^0\gamma}, \\ D^P_{\omega\omega} f^F_{\omega\pi^0\gamma} &= D^P_{\omega\omega} f^P_{\omega\pi^0\gamma} + D^P_{\omega\rho} f^P_{\rho^0\pi^0\gamma}, \end{aligned} \tag{27}$$

and hence

$$\begin{aligned} f^F_{\rho^0\pi^0\gamma} &= f^{(0)}_{\rho^0\pi^0\gamma} - \eta f^{(0)}_{\omega\pi^0\gamma} + \frac{T}{m^2_\rho - m^2_\omega + i m_\omega \Gamma_\omega} f^{(0)}_{\omega\pi^0\gamma}, \\ f^F_{\omega\pi^0\gamma} &= f^{(0)}_{\omega\pi^0\gamma} + \eta f^{(0)}_{\rho^0\pi^0\gamma} + \frac{T}{m^2_\omega - m^2_\rho + i m_\rho \Gamma_\rho} f^{(0)}_{\rho\pi^0\gamma}, \end{aligned} \tag{28}$$

where eqs.(20) and (22) have been used. Considering $|\eta| \simeq 0.006 << 1$ and $T \simeq \Pi_{\rho\omega}$, we finally obtain the desired results in the expt-MP formalism as follows

$$\begin{aligned} f_{\rho^0\pi^0\gamma}|_{\rm expt-MP} &\equiv f^F_{\rho^0\pi^0\gamma} \simeq f^{(0)}_{\rho^0\pi^0\gamma} + \frac{\Pi_{\rho\omega}}{m^2_\rho - m^2_\omega + i m_\omega \Gamma_\omega} f^{(0)}_{\omega\pi^0\gamma}, \\ f_{\omega\pi^0\gamma}|_{\rm expt-MP} &\equiv f^F_{\omega\pi^0\gamma} \simeq f^{(0)}_{\omega\pi^0\gamma} + \frac{\Pi_{\rho\omega}}{m^2_\omega - m^2_\rho + i m_\rho \Gamma_\rho} f^{(0)}_{\rho^0\pi^0\gamma}. \end{aligned} \tag{29}$$

Noting $f^{(0)}_{\omega\pi^0\gamma} = 3 f^{(0)}_{\rho^0\pi^0\gamma}$, above equations are same as eq.(1). Consequently, this result indicates that the charge-asymmetry enhancement effect to $\rho^0 \to \pi\gamma$ revealed in ref. [4] by using the effective Lagrangian theory has been confirmed in expt-PM approach.

To conclude, the Mixed Propagator (MP) approach to $\rho - \omega$ mixing is investigated in this letter. It is found that under the pole-approximation assumption the results of MP approach is not compatible both with the effective Lagrangian theory and with the experiment measurement criterion. This fact indicates the pole approximation for determining the physical basis of ρ and ω is inadequate and ad hoc. To cure these diseases, we propose a new MP approach in which the physical states of ρ and ω are determined by the requirement of experimental measurement to meson resonance. In terms of this new MP approach, the EM pion form factor F_π and form factors of $\rho^0 \to \pi^0\gamma$ and of $\omega \to \pi^0\gamma$ are derived. The results of F_π are in good agreement with data. The form factor of $\rho^0 \to \pi^0\gamma$ exhibits a hidden charge-asymmetry enhancement effect which agree with the prediction of the effective Lagrangian theory, i.e., the conclusion of ref. [4] has been confirmed.

We are pleased to acknowledge K.F.Liu for helpful discussion.

---

[1] S.Glashow, Phys.Rev.Lett., **7**, 469 (1961).

[2] H.B.O'Connell, B.C.Pearce, A.W.Thomas and A.G.Williams, Prog.Nucl.Part.Phys., **39**, 201 (1997).

[3] F.M.Renard, Springer Tracts in Modern Phys., **63**, 98-120, Springer Verlag (1972).

[4] X.J.Wang, J.H.Jiang and M.L.Yan,*Isospin Symmetry Breaking in* $\rho \to \pi\gamma$, hep-ph/0101166.

[5] M.Benayoun, S.I.Eidelman and V.N.Ivanchenko, Z.Phys., **C72**, 221 (1996).

[6] X.J.Wang and M.L.Yan, hep-ph/0010215; T.L.Zhuang, X.J.Wang and M.L.Yan, Phys. Rev., **D62**, 053007 (2000); D.N.Gao and M.L.Yan, Euro Phys Jour., **A3**, 293(1998) ; B.A.Li, Phys Rev., **D52**, 5156(1995); E.Jenkins, A.V. Manohar and M.B. Wise, Phys. Rev. Lett. **75**, 2272(1995) ; A.K. Leibovich, A.V. Manohar and M.B. Wise, Phys. Lett. **B358**, 347(1995) ; J. Bijnens, P.Gosdzinsky and P. Talavera, Phys. Lett. **B429**, 111(1998) ; J.Bijinens, P. Gosdzinsky and P.Talavera, JHEP **9801**, 014(1998) ; T.Hatsuda and T.Kunihiro, Phys. Rep. **247**, 221(1994) ; M.Bando,T.Kugo and K.Yamawaki, Nucl. Phys. **B259**, 493(1985) , Prog. Theor. Phys. **79**, 1140(1988) , Phys. Rep. **164**, 217(1988) ; N.Kaiser and U.G.Meissner, Nucl. Phys., **A519**, 671(1990) . G.Eicker, J.Gasser, A.Pich and E.de Rafel, Nucl. Phys., **B321**, 311(1989) ; G.Eicker, H.Leutwyler, J.Gasser, A.Pich and E.de Rafel, Phys. Lett., **B233**, 425(1989) ; M.Brise, Z.Phys., **355**, 231(1996) ; J.Bijinens, P.Gosdzinsky and P.Talavera, Nucl. Phys., **B501**, 495(1997) ; J.Bijnens, Phys. Rep. **265**, 369(1996).

[7] X.J. Wang and M.L. Yan, Phys. Rev. **D 62**, 094013 (2000).

[8] e.g., see: M.E.Peskin and D.V.Schroeder, *An Introduction to Quantum Field Theory*, Addison-Wesley Pub. Com., (1995), pp237.

[9] S.Gardner and H.B.O'Connell, Phys Rev., **D57** 2716 (1998).

[10] H.B.O'Connell,B.C.Pearce,A.W.Thinas,A.G.Williams,A.G.: Phys. Lett. **B336**,1 (1994) .

[11] M.Benayoun, S.Eidelman, K.Maltman, H.B.O'Connell, B.Shwartz and A.G.Williams, Europ. Phys. Jour., **C2**, 269 (1998); K.Maltman, H.B.O'Connell and A.G.Williams, Phys. Lett., **B376**, 19 (1996); S.Gardner and H.B.O'Connell, Phys. Rev., **D57**, 2716 (1998).